

The 4 Quantum Numbers

Principle Quantum Number	Secondary Quantum Number	Magnetic Quantum Number	Spin Quantum Number
n	l	m _l	m _s

Electron Configurations

Electrons fill orbitals from **lowest to highest** energy. Therefore, orbital 1s fills before 2s and 2p. However, an orbit does not necessarily fill completely before the next begins.

Types of Bonds

IONIC

- metals give electrons to non-metals
- metals form cations (+)
- non-metals form anions (-)
- this gives both atoms a stable electron configuration
- the energy level of each atom is decreased

If attraction outweighs repulsion, then a bond will form

Characteristics

- conductive in the dissolved or molten state
- solid, hard, brittle
- **high** melting point, **low** boiling point

COVALENT BONDING

1. Non-polar ▶ equal sharing of electrons for bonds
2. Polar ▶ unequal sharing of electrons, atom with higher ▲ EN is slightly +, lower ▲ EN is slightly -
3. Coordinate Covalent Bonds ▶ - both electrons forming the bond come from the same atom

Characteristics

- generally low boiling points
- solid, liquid, gas
- do not conduct electricity
- dull
- don't dissolve in water

Intermolecular Forces (cont)

3. **Dipole-Dipole** → the attraction between oppositely charged dipoles of 2 polar molecules
 - strength depends on the polarity of the molecule (more polar=stronger dipole force)
 - H-bonding is a special type which is the strongest (5% of covalent bond strength)
 - H bonded to N, O, F
 - a lone pair of electrons must be on the neighbouring molecule for the H to bond with
 - strength depends on the number of H bonds

4. **Dipole-Induced Dipole** → - nonpolar molecule forced into polarity

4. **Induced Dipole-Induced Dipole** → a.k.a. London Dispersion Forces
 - the random motion of electrons creates a temporary dipole in one nonpolar molecule. This induces polarity in the neighbouring molecule. Strength depends on # of electrons (and protons) in a molecule.

Types of Solids

- Metallic Crystals** (Metallic Bonding)
- valence electrons from a mobile sea of electrons which comprise the metallic bond
 - high melting and boiling points

- Ionic Crystals** (Ionic Bonding)
- attraction of charged ions for one another. Lattice energy is a measure of ionic strength
 - high melting and boiling points

- Covalent Crystals** (Network Covalent Bonding)
- network solids are extremely hard compounds with very high melting and boiling points due to their endless 3-D network of covalent bonds

Types of Solids (cont)

- Molecular Crystals**
- a) H-bonding
 - b) LDF
 - c) Dipole-Dipole Forces
- a) H-bonds are weaker than covalent bonds, but stronger than b) or c) below
- b) universal force of attraction between instantaneous dipoles. These forces are weak for small, low-molecular weight molecules, but large for heavy, long/highly polarizable molecules. They are stronger than c) below
- c) these forces act between **polar** molecules. They are much weaker than H-bonding

- Atomic Crystals** (Dispersion Forces)

- see section b) above

Physical properties depend on these forces.

The **stronger** the forces between particles,

- the ▲ the melting and boiling point
- the ▼ the vapour pressure
- the ▲ the viscosity
- the **greater** the surface tension

Intermolecular Forces

An attraction holding neighbouring molecules or ions together. **These are not bonds**

1. **Ion-Ion** → whole charges attract

2. **Ion-Dipole** → an ion is attracted to a polar molecule. The cation is attracted to the slightly negative portion of polar molecules and the anion to the slightly positive end



By **corinne_montpetit**
cheatography.com/corinne-montpetit/

Not published yet.
Last updated 13th October, 2017.
Page 1 of 2.

Sponsored by **CrosswordCheats.com**
Learn to solve cryptic crosswords!
<http://crosswordcheats.com>